

Methenolone Enanthate 150mg/ml

1 VIAL containing 10 ml

For multidose use

Therapeutic Class:

Anabolic Agent

Composition:

Each 1 ml of PRIMOTREX150 contains:

- 150 mg Methenolone Enanthate
- Ethyl Oleate q. s.

Indication & Dosage:

Intramuscular

Postmenopausal osteoporosis

*Adult: As enanthate: 100 mg every 2 wk,
then reduce to once every 3-4 wk after an initial response.*

Progressive breast cancer

*Adult: As enanthate: 100 mg every 1-2 wk
or 200 mg every 2-3 wk.*

Contraindications:

Hypercalcaemia or hypercalciuria, males with breast or prostate carcinoma.
Pregnancy and lactation.

Special Precautions

Lactation, CV disorders, renal or hepatic impairment, epilepsy, migraine, DM, conditions aggravated by fluid retention. Children. Regularly examine prostate in men.

Adverse Drug Reactions

Fluid and electrolyte retention, psychiatric effects, hypercalcaemia, impaired glucose tolerance, increased bone growth and skeletal weight.

Men: Impotence, priapism, azoospermia, prostatic hypertrophy; gynaecomastia, hirsutism, male pattern baldness, acne, seborrhoea, oedema, cholestatic jaundice.

Women: Veneralisation, amenorrhoea, menstrual irregularities, suppressed lactation, increased libido.

Children: Premature closure of epiphyses, stops linear growth, veneralisation symptoms.

Drug Interactions

May enhance effects of antidiabetics, ciclosporin, levothyroxine, warfarin.

Resistance to the effects of neuromuscular blockers may occur.

Lab Interference

May interfere with glucose tolerance and thyroid function tests.

Pregnancy Category

Category X: Studies in animals or human beings have demonstrated foetal abnormalities or there is evidence of foetal risk based on human experience or both, and the risk of the use of the drug in pregnant women clearly outweighs any possible benefit. The drug is contraindicated in women who are or may become pregnant.

Mechanism of Action

Methenolone is an anabolic steroid, which in high doses stimulates bone marrow functions, particularly erythropoiesis.

CIMS Class: Anabolic Agents / Cytotoxic Chemotherapy

ATC Classification: A14AA04 - metenolone ; Belongs to the class of androstan derivative anabolic steroids used as systemic anabolic agents.

Storage:

Store at room temperature, between 59 and 86 degrees F (15 and 30 degrees C), in a tightly-closed container. Store away from heat, moisture, and light. Do not store in the bathroom. Keep out of the reach of children and away from pets.

Concentrex Laboratories N. V., Moerstraat 58, 9031, Gent, Belgium

Reg. No. Licence: 253/55-C9/2010/BE

Manufactured under WHO cGMP guidelines

www.concentrexlabs.com

Date of Preparation : 19 July 2010